



Stahist[®] ***TP***

Antihistamine • Nasal Decongestant

NDC: 58407-010-10

Stahist TP[®] is a decongestant (phenylephrine) & antihistamine (triprolidine) for the treatment of allergy & upper respiratory symptoms including runny nose, sneezing, itching of the nose or throat, itchy or watery eyes, & nasal congestion. ***Stahist TP***[®] aids in reducing swelling of the nasal passages.

Stahist TP[®] bottles contain 100 tablets. Tablets are scored to facilitate half-tablet, customized dosing.

Phenylephrine HCl...10mg
Triprolidine HCl.....2.5mg



**Scan to
order!
or visit**

www.magnaweb.com

888-206-5525



DOSAGE:

Adults & Children Age 12 & Up: 1 tablet by mouth every 4 hours, not to exceed 4 tablets in 24 hours, or as directed by a physician.

Children 6 to 12 Years of Age: half a tablet by mouth every 4 hours, not to exceed 2 whole tablets in 24 hours, or as directed by a physician. *For children under 6, consult a physician.*

DRUG FACTS:

Active Ingredients: Phenylephrine HCl 10mg (nasal decongestant), Triprolidine HCl 2.5mg (antihistamine)

Uses: Temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other respiratory allergies: runny nose, sneezing, itchy nose/throat, itchy watery eyes, nasal congestion, reduces swelling of nasal passages.

Warnings: Do not exceed recommended dosage. Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis, heart disease, high blood pressure, thyroid disease, diabetes, glaucoma or trouble urinating due to an enlarged prostate gland. Ask a doctor before use if you are taking sedatives or tranquilizers. When using this product, excitability may occur, especially in children. May cause drowsiness. Alcohol, sedatives and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages. Use caution when driving a motor vehicle or operating machinery. Stop use and ask a doctor if nervousness, dizziness, sleeplessness or new symptoms occur. Cough or nasal congestion persists for more than one week, tends to recur or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition. If pregnant or breast-feeding, ask a healthcare professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.