

Ahist™

Antihistamine

- The antihistamine chlorcyclizine HCl has a prolonged action and a lower incidence of drowsiness (Hardman, Limbird & Gilman, 2006).
- Better anticholinergic control of rhinorrhea [and postnasal drip] among first-generation antihistamines (Wallace & Dykewicz, 2008).
- Dosage can be titrated up to three times per day (up to one tablet every 8 hours, maximum three tablets per day) according to individual patient needs; tablet is scored.
- Low cost, great value!

Dosage:

Adults and Children over 12 years of age: 1 tablet by mouth every 6-8 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor.

Children 6 to 12 years of age: ½ tablet every 6-8 hours, not to exceed 1½ tablets in 24 hours, or as directed by a doctor.

Children under 6 years of age: consult a physician.

Ahist™

NDC #58407-025-30
Antihistamine

Drug Facts

Active Ingredients (In each immediate-release tablet)

Chlorcyclizine HCl 25 mg - Antihistamine

Uses

Temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Warnings Do not exceed recommended dosage.

Do not take this product unless directed by a doctor if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers.

When using this product:

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness effect
- use caution when driving a motor vehicle or operating machinery.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions: Do not exceed recommended dosage.

Adults and children 12 years of age and over: 1 tablet by mouth every 6 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor

Children 6 to under 12 years of age: ½ tablet by mouth every 6 hours, not to exceed 1½ tablets in 24 hours, or as directed by a doctor

Children under 6 years of age Consult a physician

Store at 59° - 86°F (15° - 30°C) [see USP Controlled Room Temperature]

Inactive ingredients: Lake Blend Green, Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate

Questions or Comments?

Manufactured for:

MAGNA
Pharmaceuticals, Inc.
...commitment to...
...responsibility...

Louisville, KY 40299
magnaweb.com



The statements in this brochure have not been evaluate by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.