

Stahist[®] TP

Antihistamine • Nasal Decongestant

Each **Stahist[®] TP** tablet contains:

Phenylephrine HCl 10mg
Triprolidine HCl 2.5mg

Dosage: Adults and children 12 years of age and over:

1 tablet by mouth every 4 hours, not to exceed 4 tablets in 24 hours, or as directed by a physician.

Children 6 to 12 years of age: 1/2 tablet every 4 hours, not to exceed 2 tablets in 24 hours, or as directed by a physician. **Under 6:** Consult a physician.



- **Tried & true drying formula.**
- **Small scored tablet for customized dosing.**
- **Relieves runny nose, sneezing, itchy nose/throat, itchy/watery eyes, congestion, swollen nasal passages.**

See Drug Facts on Back
NDC 58407-0010-10
MAGNAWEB.COM

Stahist TP[®]

NDC 58407-0010-10

Antihistamine • Nasal Decongestant

Drug Facts

Active Ingredients (in each tablet)

Phenylephrine HCl 10 mg - Nasal Decongestant

Triprolidine HCl 2.5 mg - Antihistamine

Uses

temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ reduces swelling of nasal passages

Warnings

■ Do not exceed recommended dosage.

Do not use this product

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

■ excitability may occur, especially in children ■ may cause drowsiness ■ alcohol, sedatives and tranquilizers may increase the drowsiness effect ■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur ■ cough or nasal congestion persists for more than 1 week, tends to recur or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition. ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison

Control Center right away.

Directions: Do not exceed recommended dosage.

Adults and Children 12 years of age and over: 1 tablet by mouth every 4 hours, not to exceed 4 tablets in 24 hours, or as directed by a doctor.

Children 6 to under 12 years of age: 1/2 tablet every 4 hours, not to exceed 2 tablets in 24 hours, or as directed by a doctor.

Children under 6 years of age: Consult a doctor

Other Information

Store at 15°-30° C (59°-86° F)

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Inactive ingredients

Magnesium stearate, microcrystalline cellulose, sodium starch glycolate

Question or comments?

Call 1-888-206-5525

Manufactured for:

MAGNA
Pharmaceuticals, Inc.
— Commitment to —

Louisville, KY 40299
magnaweb.com